

# HILBERT COEFFICIENTS AND DEPTH OF FIBER CONES

A. V. JAYANTHAN AND J. K. VERMA

*Dedicated to Prof. Wolmer Vasconcelos on the occasion of his 65th birthday*

**ABSTRACT.** Criteria are given in terms of certain Hilbert coefficients for the fiber cone  $F(I)$  of an  $\mathfrak{m}$ -primary ideal  $I$  in a Cohen-Macaulay local ring  $(R, \mathfrak{m})$  so that it is Cohen-Macaulay or has depth at least  $\dim(R) - 1$ . A version of Huneke's fundamental lemma is proved for fiber cones. Goto's results concerning Cohen-Macaulay fiber cones of ideals with minimal multiplicity are obtained as consequences.

## 1. INTRODUCTION

Let  $(R, \mathfrak{m})$  be a  $d$ -dimensional Cohen-Macaulay local ring having infinite residue field  $R/\mathfrak{m}$ . Let  $I$  be an  $\mathfrak{m}$ -primary ideal. The objective of this paper is to explore some connections between the coefficients of the polynomial  $P_{\mathfrak{m}}(I, n)$  corresponding to the function  $H_{\mathfrak{m}}(I, n) = \lambda(R/\mathfrak{m}I^n)$  and depth of the fiber cone  $F(I) = \bigoplus_{n \geq 0} I^n/\mathfrak{m}I^n$  of  $I$ . The relation between Hilbert coefficients and depth has been a subject of several papers in the context of associated graded rings and Rees algebras of ideals. That conditions on Hilbert coefficients could force high depth for associated graded rings was first observed by Sally in [S1]. Since then numerous conditions have been provided for the Hilbert coefficients so that the associated graded ring of  $I$ ,  $G(I) = \bigoplus_{n \geq 0} I^n/I^{n+1}$ , is either Cohen-Macaulay or has almost maximal depth, i.e. the grade of the maximal homogeneous ideal of  $G(I)$  is at least  $d - 1$ .

Let  $J$  be a minimal reduction of  $I$ . In their elegant paper [HM] Huckaba and Marley provided necessary and sufficient conditions on the coefficients of the Hilbert polynomial  $P(I, n)$  corresponding to the Hilbert function  $H(I, n) = \lambda(R/I^n)$  so that  $G(I)$  is Cohen-Macaulay and has almost maximal depth. We shall write the Hilbert polynomial  $P(I, n)$  in the following way:

$$P(I, n) = e_0(I) \binom{n+d-1}{d} - e_1(I) \binom{n+d-2}{d-1} + \cdots + (-1)^d e_d.$$

Huckaba and Marley showed:

- (i)  $G(I)$  is Cohen-Macaulay if and only if  $e_1(I) = \sum_{n \geq 1} \lambda(I^n + J/J)$
- (ii)  $G(I)$  has almost maximal depth if and only if  $e_1(I) = \sum_{n \geq 1} \lambda(I^n/JI^{n-1})$ .

Their results unify several results known in the literature and provide an effective approach to deal with such questions.

The relation between Hilbert coefficients and depth of  $F(I)$  is not well understood. The papers [CZ], [G], [DV] and [DRV] provide sufficient conditions in terms of certain Hilbert coefficients for the

---

The first author was supported by the National Board for Higher Mathematics, India.

AMS Classification 2000: 13H10, 13H15 (Primary), 13C15, 13A02 (Secondary).

Cohen-Macaulay property of  $F(I)$ . Recently, a number of papers have appeared which discuss Cohen-Macaulay and other properties of fiber cones. See for example [CGPU], [C], [CPV], [HH]. The form ring  $G(I)$  and the fiber cone  $F(I)$  coincide when  $I = \mathfrak{m}$ . This indicates that there may be analogues of results of Huckaba and Marley for the fiber cone. The first guess for the appropriate Hilbert function to be used is naturally the Hilbert function of  $F(I)$ . However we have observed that this does not seem to be of much help in predicting depth. We will show that the coefficients of the polynomial  $P_{\mathfrak{m}}(I, n)$  corresponding to the function  $\lambda(R/\mathfrak{m}I^n)$  control the Cohen-Macaulay and almost maximal depth properties of  $F(I)$ .

We now describe the main results of this paper. Write the polynomial  $P_{\mathfrak{m}}(I, n)$  as

$$P_{\mathfrak{m}}(I, n) = g_0(I) \binom{n+d-1}{d} - g_1(I) \binom{n+d-2}{d-1} + \cdots + (-1)^d g_d(I).$$

Let  $\text{grade } G(I)_+ \geq d-1$ . In sections 4 and 5 we shall prove that

- (i)  $F(I)$  is Cohen-Macaulay if and only if  $g_1(I) = \sum_{n \geq 1} \lambda(\mathfrak{m}I^n + JI^{n-1}/JI^{n-1}) - 1$  and
- (ii)  $F(I)$  has almost maximal depth if and only if  $g_1(I) = \sum_{n \geq 1} \lambda(\mathfrak{m}I^n/\mathfrak{m}JI^{n-1}) - 1$ .

It can be seen that the minimal number of generators of  $I$ ,  $\mu(I) \leq e_0(I) + d - \lambda(R/I)$ . Goto in [G] defined an ideal  $I$  in a Cohen-Macaulay local ring to have *minimal multiplicity* if equality holds in the above inequality. He showed that if  $I$  has minimal multiplicity then  $F(I)$  is Cohen-Macaulay if and only if  $G(I)$  has almost maximal depth. We shall recover this result in section 6 as a consequence of our criterion for Cohen-Macaulayness in terms of  $g_1(I)$ . In fact we shall prove that  $I$  has minimal multiplicity if and only if  $g_1(I) = -1$ . We will also recover a bound for the multiplicity of the fiber cone due to Corso, Polini and Vasconcelos [CPV].

Since the criteria for Cohen-Macaulay and almost maximal depth properties of  $F(I)$  require one to know the coefficient  $g_1(I)$ , it is desirable to have an effective method of its computation. In Section 5 we shall show that in a one dimensional Cohen-Macaulay local ring  $g_1(I) = \sum_{n \geq 1} \lambda(\mathfrak{m}I^n/J\mathfrak{m}I^{n-1}) - 1$ . We will also present a simple proof of a criterion due to Cortadellas and Zarzuela [CZ] for the sequence of initial forms in  $F(I)$  of elements in a regular sequence in  $R$  to be a regular sequence in  $F(I)$ .

In Section 3 we shall generalize the *fundamental lemma of Huneke* in [H] for finding a formula for  $g_1(I)$ . However, we need a modified version of this lemma so that it works for the function  $\lambda(R/\mathfrak{m}I^n)$ .

In the second section we will discuss the technical topic of superficial and regular elements in fiber cones.

As no extra effort is required, we will prove our results for filtrations of ideals. In a subsequent paper on fiber cones we will see that it is useful to develop the criteria for filtrations as sometime we need to deal with filtrations to prove results about the  $I$ -adic filtration.

*Acknowledgements:* We thank Balwant Singh and Santiago Zarzuela for discussions.

## 2. SUPERFICIAL AND REGULAR ELEMENTS IN FIBER CONES OF FILTRATIONS

In this section we will gather some results on superficial and regular elements in fiber cones. Throughout this paper  $(R, \mathfrak{m})$  will denote a Noetherian local ring of positive (Krull) dimension  $d$ , with maximal ideal  $\mathfrak{m}$  and infinite residue field  $R/\mathfrak{m}$ . A sequence of ideals  $\mathcal{F} = \{I_n\}_{n \geq 0}$  is called a

filtration if  $I_0 = R$ ,  $I_1 \neq R$ ,  $I_{n+1} \subseteq I_n$ , and  $I_n I_m \subseteq I_{n+m}$  for all  $n, m \geq 0$ . The Rees algebra  $\mathcal{R}(\mathcal{F})$  and the associated graded ring  $G(\mathcal{F})$  are defined to be the graded rings:

$$\mathcal{R}(\mathcal{F}) = R \oplus I_1 t \oplus I_2 t^2 \oplus \cdots, G(\mathcal{F}) = R/I_1 \oplus I_1/I_2 \oplus I_2/I_3 \oplus \cdots$$

The filtration  $\mathcal{F}$  is called Noetherian if  $\mathcal{R}(\mathcal{F})$  is a Noetherian ring. Throughout the paper we will assume that  $\mathcal{F}$  is Noetherian and  $I_n \neq 0$  for all  $n \geq 0$ . The ideal generated by elements of positive degree in  $G(\mathcal{F})$  will be denoted by  $G(\mathcal{F})_+$ . The filtration  $\mathcal{F}$  is called  $I_1$ -good if  $\mathcal{R}(\mathcal{F})$  is a finite module over the Rees algebra  $\mathcal{R}(I_1)$ . An  $I_1$ -good filtration is called a Hilbert filtration if  $I_1$  is  $\mathfrak{m}$ -primary.

The fiber cone of  $\mathcal{F}$  with respect to an ideal  $K$  containing  $I_1$  is the graded ring

$$F_K(\mathcal{F}) = R/K \oplus I_1/KI_1 \oplus I_2/KI_2 \oplus \cdots.$$

For  $x \in I_1$ , let  $x^*$  and  $x^o$  denote the initial form in degree one component of  $G(\mathcal{F})$  and  $F_K(\mathcal{F})$  respectively. We will always assume that  $I_{n+1} \subseteq KI_n$  for all  $n \geq 0$ . This is required in Lemma 2.3 which is essential in all the arguments that apply induction on the dimension of  $R$  in the subsequent sections.

**Definition 2.1.** For an element  $x \in I_1$  such that  $x^o \neq 0 \in F_K(\mathcal{F})$ ,  $x^o$  is said to be superficial in  $F_K(\mathcal{F})$  if there exists  $c > 0$  such that  $(0 : x^o) \cap F_K(\mathcal{F})_n = 0$  for all  $n > c$ .

It can easily be seen that  $x^o$  is superficial in  $F_K(\mathcal{F})$  if and only if there exists  $c > 0$  such that  $(KI_{n+1} : x) \cap I_n = KI_n$  for all  $n > c$ . We first show the existence of superficial elements in  $F_K(\mathcal{F})$  and proceed to prove some of their properties. The existence of superficial elements in a graded ring is well-known. Since in our case we need existence of elements which are superficial in  $F_K(\mathcal{F})$  as well as  $G(\mathcal{F})$  simultaneously, we give a proof of the following result for the sake of completeness.

**Proposition 2.2.** Let  $(R, \mathfrak{m})$  be a Noetherian local ring of dimension  $d > 0$ . Let  $\mathcal{F}$  be an  $I_1$ -good filtration and  $K$  be an  $\mathfrak{m}$ -primary ideal containing  $I_1$ . Then there exists  $x \in I_1 \setminus KI_1$  such that  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ .

*Proof.* Let the set of associated primes of  $G(\mathcal{F})$  and  $F_K(\mathcal{F})$  be

$$\text{Ass}(G(\mathcal{F})) = \{P_1, \dots, P_r, P_{r+1}, \dots, P_s\} \text{ and } \text{Ass}(F_K(\mathcal{F})) = \{Q_1, \dots, Q_l, Q_{l+1}, \dots, Q_m\}$$

such that for all  $n \gg 0$ ,  $I_n/I_{n+1} \subseteq P_i$  for  $r+1 \leq i \leq s$  and  $I_n/KI_n \subseteq Q_j$  for  $l+1 \leq j \leq m$ . The associated graded ring  $G(\mathcal{F})$  and the fiber cone  $F_K(\mathcal{F})$  are both homomorphic images of the extended Rees algebra  $\mathcal{R}(\mathcal{F})(t^{-1})$  since  $G(\mathcal{F}) = \mathcal{R}(\mathcal{F})(t^{-1})/(t^{-1})$  and  $F_K(\mathcal{F}) = \mathcal{R}(\mathcal{F})(t^{-1})/(t^{-1}, K)$ . Let  $\mathcal{P} = \{P'_1, \dots, P'_s, Q'_1, \dots, Q'_m\}$  be the collection of prime ideals in the extended Rees algebra  $\mathcal{R}(\mathcal{F})(t^{-1})$  which are the pre-images of the ideals  $P_1, \dots, P_s$  in  $\text{Ass}(G(\mathcal{F}))$  and  $Q_1, \dots, Q_m$  in  $\text{Ass}(F_K(\mathcal{F}))$  respectively. Since  $R/\mathfrak{m}$  is infinite,  $\mathcal{R}_1 \neq \mathfrak{m}I_1 t \cup_{i=1}^r P'_i \cup_{i=1}^l Q'_i$ . Choose  $xt \in \mathcal{R}_1 \setminus \mathfrak{m}I_1 t \cup_{i=1}^r P'_i \cup_{i=1}^l Q'_i$ . We show that  $0 \neq x^o \in F_K(\mathcal{F})_1$  is superficial in  $F_K(\mathcal{F})$  and  $x^* \in G(\mathcal{F})_1$  is superficial in  $G(\mathcal{F})$ . We need to show that  $(0 : x^o) \cap F_K(\mathcal{F})_n = 0$  for  $n \gg 0$ . Let  $y^o \in (0 : x^o)$ . Let  $(0) = N_1 \cap \cdots \cap N_m$  be a primary decomposition of  $(0)$  in  $F_K(\mathcal{F})$  such that  $N_i$  is  $Q_i$ -primary for  $i = 1, \dots, m$ . Then  $y^o x^o \in N_i$  for all  $1 \leq i \leq l$  and  $x^o \notin Q_i$  for  $i = 1, \dots, l$ . Therefore  $y^o \in N_i$  for  $i = 1, \dots, l$ . Thus

$(0 : x^o) \subseteq N_1 \cap \cdots \cap N_l$ . For  $l+1 \leq j \leq m$ ,  $F_K(\mathcal{F})_n \subseteq Q_j$  for  $n \gg 0$ . Therefore there exists a  $c > 0$  such that  $\oplus_{n \geq c} F_K(\mathcal{F})_n \subseteq N_{l+1} \cap \cdots \cap N_m$ . Therefore for all  $n \geq c$

$$(0 : x^o) \cap F_K(\mathcal{F})_n \subseteq N_1 \cap \cdots \cap N_l \cap N_{l+1} \cap \cdots \cap N_m = (0).$$

Therefore  $x^o$  is superficial in  $F_K(\mathcal{F})$ . A similar argument shows that  $x^*$  is superficial in  $G(\mathcal{F})$ .  $\square$

In the next lemma, we characterize the property of an element being superficial in the fiber cone in terms of its properties in the local ring.

**Lemma 2.3.** (i) *If there exists a  $c > 0$  such that  $(KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ , then  $x^o$  is superficial in  $F_K(\mathcal{F})$ .*

(ii) *If  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ , then there exists  $c > 0$  such that  $(KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ . Moreover if  $x$  is regular in  $R$ , then  $KI_n : x = KI_{n-1}$  for all  $n \gg 0$ .*

*Proof.* (i) Suppose  $(KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ . Then  $(KI_n : x) \cap I_{n-1} \subseteq (KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ . Therefore  $x^o$  is superficial in  $F_K(\mathcal{F})$ .

(ii) Suppose  $x \in I_1$  is such that  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ . Then there exist  $c_1, c_2$  such that for all  $n > c_1$ ,  $(0 : x^*) \cap G(\mathcal{F})_n = 0$  and  $(0 : x^o) \cap F_K(\mathcal{F})_n = 0$  for all  $n > c_2$ . Choose  $c = \max\{c_1, c_2\} + 1$ .

*Claim :*  $(KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ .

Suppose that there exists an integer  $t$  such that  $y \in I_t \setminus KI_t$ . If  $xy \in KI_{t+1}$ , then  $y^o \in (0 : x^o) \cap F_K(\mathcal{F})_t = 0$  which is a contradiction (the last equality holds because  $t \geq c > c_2$ ). Therefore,  $xy \notin KI_{t+1}$ . Since  $xy \in KI_n$ ,  $n < t+1$  so that  $n \leq t$ . Therefore  $y \in I_t \subseteq I_n \subseteq KI_{n-1}$ .

Now, suppose that the image of  $y$  in  $F_K(\mathcal{F})$  is zero, i.e., given any  $n$  such that  $y \in I_n$ ,  $y \in KI_n$ . Then  $y \in (KI_n : x) \cap I_c \subseteq (I_n : x) \cap I_c = I_{n-1}$ , since  $c > c_1$ . Therefore,  $y \in KI_{n-1}$ .

Suppose that  $x$  is regular in  $R$ . Then by the Artin-Rees lemma, there exists a  $c$  such that  $KI_n \cap (x) = I_{n-c}(KI_c \cap (x)) \subseteq xI_{n-c} \subseteq xI_c$  for all  $n \geq 2c$ . Therefore  $KI_n : x \subseteq I_c$  for  $n \gg 0$ . Hence for all  $n \gg 0$ ,  $KI_n : x = (KI_n : x) \cap I_c = KI_{n-1}$ .  $\square$

For an element  $x \in I$  such that  $x^*$  is superficial in  $G(I)$ , it is known that  $x \in I \setminus I^2$ . In the following result we show that a similar property is true for superficial elements in  $F_K(\mathcal{F})$ .

**Lemma 2.4.** *If  $x^o \in I_1/KI_1$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ , then  $x \in I_1 \setminus KI_1$ .*

*Proof.* Since  $x^o \in F_K(\mathcal{F})$  and  $x^* \in G(\mathcal{F})$  are superficial, by Lemma 2.3, there exists  $c > 0$  such that  $(KI_n : x) \cap I_c = KI_{n-1}$  for all  $n > c$ . Put  $n = c+1$ . Then  $(KI_{c+1} : x) \cap I_c = KI_c$ . Suppose  $x \in KI_1$ . Let  $y \in I_c$ . Then  $xy \in KI_{c+1}$  so that  $y \in (KI_{c+1} : x) \cap I_c = KI_c$ . Therefore  $I_c = KI_c$ . By Nakayama Lemma  $I_c = 0$  which is a contradiction to the fact that  $\mathcal{F}$  is a Hilbert filtration. Therefore  $x \in I_1 \setminus KI_1$ .  $\square$

For the fiber cone  $F_K(\mathcal{F})$ , let  $H(F_K(\mathcal{F}), n) = \lambda(F_K(\mathcal{F})_n) = \lambda(I_n/KI_n)$  denote its Hilbert function and let  $P(F_K(\mathcal{F}), n)$  denote the corresponding Hilbert polynomial.

**Proposition 2.5.** *Let  $x^o \in F_K(\mathcal{F})$  be superficial. Then  $\dim F_K(\mathcal{F})/x^o F_K(\mathcal{F}) = \dim F_K(\mathcal{F}) - 1$ .*

*Proof.* Consider the exact sequence

$$0 \longrightarrow (KI_n : x) \cap I_{n-1}/KI_{n-1} \longrightarrow I_{n-1}/KI_{n-1} \xrightarrow{x} I_n/KI_n \longrightarrow I_n/(KI_n + xI_{n-1}) \longrightarrow 0.$$

Then  $H(F_K(\mathcal{F}), n) - H(F_K(\mathcal{F}), n-1) = H(F_K(\mathcal{F})/x^o F_K(\mathcal{F}), n) - \lambda((KI_n : x) \cap I_{n-1}/KI_{n-1})$  for all  $n \geq 1$ . Since  $x^o$  is superficial in  $F_K(\mathcal{F})$ ,  $(KI_n : x) \cap I_{n-1} = KI_{n-1}$  for  $n \gg 0$ , so that  $P(F_K(\mathcal{F})/(x^o), n) = P(F_K(\mathcal{F}), n) - P(F_K(\mathcal{F}), n-1)$ . Hence  $\dim F_K(\mathcal{F})/x^o F_K(\mathcal{F}) = \dim F_K(\mathcal{F}) - 1$ .  $\square$

In the following lemma we provide a characterization for regular elements in  $F_K(\mathcal{F})$ . It can be seen that this property is quite similar to the behaviour of regular elements in  $G(I)$ .

**Lemma 2.6.** *For  $x \in I_1 \setminus KI_1$ ,  $x^o \in F_K(\mathcal{F})$  is regular if and only if  $(KI_n : x) \cap I_{n-1} = KI_{n-1}$  for all  $n \geq 1$ . If  $x^*$  is regular in  $G(\mathcal{F})$  and  $x^o$  is regular in  $F_K(\mathcal{F})$  then  $KI_n : x = KI_{n-1}$  for all  $n \geq 1$ .*

*Proof.* Suppose  $(KI_n : x) \cap I_{n-1} = KI_{n-1}$  for all  $n \geq 1$ . In other words,  $(0 : x^o) \cap F_K(\mathcal{F})_{n-1} = (0)$  for all  $n \geq 1$ . Since  $(0 : x^o)$  is a homogeneous ideal,  $(0 : x^o) = \bigoplus_{n \geq 0} (0 : x^o) \cap F_K(\mathcal{F})_n = (0)$ . Conversely, assume that  $x^o$  is a regular element in  $F_K(\mathcal{F})$ . Then  $\alpha_n : F_K(\mathcal{F})_{n-1} \longrightarrow F_K(\mathcal{F})_n$  is an injective map for all  $n \geq 1$ , where  $\alpha_n$  is the multiplication by  $x^o$  for all  $n \geq 1$ . Since  $\ker \alpha_n = (KI_n : x) \cap I_{n-1}/KI_{n-1} = 0$ ,  $(KI_n : x) \cap I_{n-1} = KI_{n-1}$  for all  $n \geq 1$ .

If  $KI_n : x = KI_{n-1}$  for all  $n \geq 1$ , then clearly  $x^o$  is regular in  $F_K(\mathcal{F})$ . Suppose that  $x^*$  is regular in  $G(\mathcal{F})$  and  $x^o$  is regular in  $F_K(\mathcal{F})$ . Let  $y \in KI_n : x$ . If there exists a  $t$  such that  $0 \neq y^o \in I_t/KI_t$ , then  $0 \neq y^o x^o \in I_{t+1}/KI_{t+1}$  so that  $yx \notin KI_{t+1}$ . Since  $yx \in KI_n$ ,  $n < t+1$ . Therefore  $y \in I_t \subseteq I_n \subseteq KI_{n-1}$ . Suppose we can not find  $t$  such that  $0 \neq y^o \in I_t/KI_t$ . Then, if  $y \in I_n$ ,  $y \in KI_n$ . Since  $y \in KI_n : x$ ,  $y \in I_n : x$ . By hypothesis,  $x^*$  is regular in  $G$ . Hence  $I_n : x = I_{n-1}$  for  $n \geq 1$ . Therefore  $y \in I_{n-1}$  and hence  $y \in KI_{n-1}$ .  $\square$

The following lemma is an analogue of Lemma 2.2 of [HM]. This will play a crucial role in induction arguments. This is the so-called Sally-machine for fiber cones.

**Lemma 2.7.** *Let  $x \in I_1$  be such that  $x^*$  is superficial in  $G(\mathcal{F})$  and  $x^o \in F_K(\mathcal{F})$  is superficial in  $F_K(\mathcal{F})$ . Let  $\bar{\mathcal{F}} = \{I_n + xR/xR\}_{n \geq 0}$  and  $\bar{K} = K/xR$ . If  $\text{depth } F_{\bar{K}}(\bar{\mathcal{F}}) > 0$ , then  $x^o$  is regular in  $F_K(\mathcal{F})$ .*

*Proof.* Let  $y^o \in I_t/KI_t$  be such that its natural image  $\bar{y}^o$  is a regular element in  $F_{\bar{K}}(\bar{\mathcal{F}})$ . Then  $(KI_{n+tj} : y^j) \cap I_n \subseteq (KI_n, x)$  for all  $n, j \geq 1$ . Since  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ , there exists  $c > 0$  such that  $(KI_{n+j} : x^j) \cap I_c = KI_n$  for all  $n > c$  and  $j \geq 1$ , by Lemma 2.3 (ii). Let  $n$  and  $j$  be arbitrary and  $p > c/t$ . Then  $y^p(KI_{n+j} : x^j) \subseteq (KI_{n+pt+j} : x^j) \cap I_c \subseteq KI_{n+pt}$ . Thus

$$(KI_{n+j} : x^j) \cap I_n \subseteq (KI_{n+pt} : y^p) \cap I_n \subseteq (KI_n, x).$$

Therefore  $(KI_{n+j} : x^j) \cap I_n \subseteq KI_n + x(KI_{n+j} : x^{j+1})$ . Iterating this formula  $n+1$  times we get,

$$\begin{aligned} (KI_{n+j} : x^j) \cap I_n &\subseteq KI_n + xKI_{n-1} + \cdots + x^{n+1}(KI_{n+j} : x^{n+j+1}) \\ &= KI_n. \end{aligned}$$

Therefore  $(KI_{n+j} : x^j) \cap I_n = KI_n$  for all  $n \geq 1$  and hence  $x^o$  is regular in  $F_K(\mathcal{F})$ .  $\square$

### 3. HILBERT COEFFICIENTS FOR THE FUNCTION $\lambda(R/KI_n)$

Throughout this section  $\mathcal{F} = \{I_n\}_{n \geq 0}$  will be a Hilbert filtration of  $R$ . Let  $K$  be an ideal of  $R$  such that  $I_{n+1} \subseteq KI_n$  for all  $n \geq 0$ . Let  $H(F, n) = \lambda(F_K(\mathcal{F})_n) = \lambda(I_n/KI_n)$  be the Hilbert function of the fiber cone  $F = F_K(\mathcal{F})$ . Then,  $H(F, n) = \lambda(R/KI_n) - \lambda(R/I_n)$ . Since both  $H(F, n)$  and  $\lambda(R/I_n)$  are polynomials for  $n \gg 0$ ,  $\lambda(R/KI_n)$  is also a polynomial for  $n \gg 0$ . Since the coefficients of this polynomial are related with the Hilbert coefficients of the fiber cone and the Hilbert-Samuel coefficients of  $\mathcal{F}$ , it is expected that their properties will be related with the properties of the fiber cone. Huneke's fundamental lemma [H] provides formulas for the Hilbert coefficients of the Hilbert polynomial of an  $\mathfrak{m}$ -primary ideal in a two-dimensional Cohen-Macaulay local ring. We will prove an analogue of this lemma for the fiber cones. It will yield formulas for the Hilbert polynomial of the fiber cone once we have access to a minimal reduction of  $I_1$ .

Let  $H_K(\mathcal{F}, n) = \lambda(R/KI_n)$  (resp.  $H(\mathcal{F}, n) = \lambda(R/I_n)$ ) and let  $P_K(\mathcal{F}, n)$  (resp.  $P(\mathcal{F}, n)$ ) be the corresponding polynomial. Since  $P_K(\mathcal{F}, n) = P(\mathcal{F}, n) + P(F, n)$ , it is a polynomial of degree  $d$  with leading coefficient  $e_0(I)$ . We write the above polynomials in the following way:

$$\begin{aligned} P(F, n) &= f_0 \binom{n+d-2}{d-1} - f_1 \binom{n+d-3}{d-2} + \cdots + (-1)^{d-1} f_{d-1}, \\ P(\mathcal{F}, n) &= e_0 \binom{n+d-1}{d} - e_1 \binom{n+d-2}{d-1} + \cdots + (-1)^d e_d, \\ P_K(\mathcal{F}, n) &= g_0 \binom{n+d-1}{d} - g_1 \binom{n+d-2}{d-1} + \cdots + (-1)^d g_d. \end{aligned}$$

Then  $g_0 = e_0$  and  $g_i = e_i - f_{i-1}$  for all  $1 \leq i \leq d$ . Note that  $P(F, n)$ , given above, is not written in the standard way of representing the Hilbert function of a graded algebra. The changed notation is for the convenience of the computation. For a numerical function  $h : \mathbb{Z} \rightarrow \mathbb{Z}$ , let  $\Delta h(n) := h(n) - h(n-1)$ .

**Lemma 3.1.** *Let  $(R, \mathfrak{m})$  be a 2-dimensional Cohen-Macaulay local ring. Let  $\mathcal{F}$  be a Hilbert filtration,  $K$  be an ideal with  $I_1 \subseteq K$  and  $J = (x, y)$  be a minimal reduction of  $I_1$ . Then for all  $n \geq 2$ ,*

$$\Delta^2 [P_K(\mathcal{F}, n) - H_K(\mathcal{F}, n)] = \lambda \left( \frac{KI_n}{KJI_{n-1}} \right) - \lambda \left( \frac{KI_{n-1} : J}{KI_{n-2}} \right).$$

*Proof.* Consider the exact sequence

$$0 \longrightarrow \frac{R}{KI_{n-1}:J} \xrightarrow{\beta} \left( \frac{R}{KI_{n-1}} \right)^2 \xrightarrow{\alpha} \frac{J}{KJI_{n-1}} \longrightarrow 0,$$

where  $\alpha$  is the map  $\alpha(\bar{r}, \bar{s}) = \overline{xr + ys}$  and  $\beta(\bar{r}) = (\bar{y}\bar{r}, -\bar{x}\bar{r})$ . It follows that for all  $n \geq 2$

$$\begin{aligned} 2\lambda(R/KI_{n-1}) &= \lambda(R/(KI_{n-1} : J)) + \lambda(J/KJI_{n-1}) \\ &= \lambda(R/(KI_{n-1} : J)) + \lambda(R/KJI_{n-1}) - \lambda(R/J). \end{aligned}$$

Therefore  $e_0(\mathcal{F}) + 2\lambda(R/KI_{n-1}) = \lambda(R/KJI_{n-1}) + \lambda(R/(KI_{n-1} : J))$ . Hence

$$\begin{aligned} e_0(\mathcal{F}) &= \lambda(R/KI_n) + 2\lambda(R/KI_{n-1}) - \lambda(R/KI_{n-2}) \\ &= \lambda(R/KJI_{n-1}) - \lambda(R/KI_n) + \lambda(R/(KI_{n-1} : J)) - \lambda(R/KI_{n-2}) \\ &= \lambda(KI_n/KJI_{n-1}) - \lambda(KI_{n-1} : J/KI_{n-2}) \end{aligned}$$

Since  $\Delta^2 P_K(\mathcal{F}, n) = e_0(\mathcal{F})$ ,

$$\Delta^2 [P_K(\mathcal{F}, n) - H_K(\mathcal{F}, n)] = \lambda \left( \frac{KI_n}{KJI_{n-1}} \right) - \lambda \left( \frac{KI_{n-1} : J}{KI_{n-2}} \right).$$

□

**Corollary 3.2** ([H], Fundamental Lemma 2.4). *Let  $(R, \mathfrak{m})$  be a 2-dimensional Cohen-Macaulay local ring and  $I$  be an  $\mathfrak{m}$ -primary ideal. Let  $J = (x, y)$  be a minimal reduction of  $I$ . Let  $H(I, n) = \lambda(R/I^n)$  be the Hilbert function of  $I$  and let  $P(I, n)$  be the corresponding Hilbert polynomial. Then for all  $n \geq 2$ ,*

$$\Delta^2 [P(I, n) - H(I, n)] = \lambda \left( \frac{I^n}{JI^{n-1}} \right) - \lambda \left( \frac{I^{n-1} : J}{I^{n-2}} \right).$$

*Proof.* Set  $K = R$ ,  $\mathcal{F} = \{I^n\}$  in Lemma 3.1. Then  $H_K(\mathcal{F}, n) = H(I, n)$  for all  $n \geq 0$  so that  $P_K(\mathcal{F}, n) = P(I, n)$  and hence the assertion follows. □

As a consequence of the generalization of the Fundamental Lemma, we obtain expressions for the Hilbert coefficients  $g_1$  and  $g_2$ .

**Corollary 3.3.** *Set*

$$v_n = \begin{cases} e_0(\mathcal{F}) & \text{if } n = 0 \\ e_0(\mathcal{F}) - \lambda(R/KI_1) + \lambda(R/K) & \text{if } n = 1 \\ \lambda(KI_n/KJI_{n-1}) - \lambda(KI_{n-1} : J/KI_{n-2}) & \text{if } n \geq 2. \end{cases}$$

*Then  $g_1 = \sum_{n \geq 1} v_n$  and  $g_2 = \sum_{n \geq 1} (n-1)v_n + \lambda(R/K)$ .*

*Proof.* From Lemma 3.1 we have,

$$\sum_{n \geq 0} \Delta^2 [P_K(\mathcal{F}, n) - H_K(\mathcal{F}, n)] t^n = \sum_{n \geq 0} v_n t^n.$$

Write  $P_K(\mathcal{F}, n) = e_0(\mathcal{F}) \binom{n+2}{2} - g'_1(n+1) + g'_2$ . Then comparing with the earlier notation, we get  $g_1 = g'_1 - e_0(\mathcal{F})$  and  $g_2 = e_0(\mathcal{F}) - g'_1 + g'_2$ . Since  $P_K(\mathcal{F}, n)$  is a polynomial of degree 2,  $\Delta^2 P_K(\mathcal{F}, n) = e_0(\mathcal{F})$  for all  $n \geq 0$  so that  $\sum_{n \geq 0} \Delta^2 P_K(\mathcal{F}, n) t^n = e_0(\mathcal{F})/(1-t)$ . Let  $\sum_{n \geq 0} H_K(\mathcal{F}, n) t^n = f(t)/(1-t)^3$ .

Then by Proposition 4.1.9 of [BH],  $e_0(\mathcal{F}) = f(1)$ ,  $g'_1 = f'(1)$  and  $g'_2 = f''(1)/2!$ . Also we have,

$$\begin{aligned}
\sum_{n \geq 0} \Delta^2 H_K(\mathcal{F}, n) t^n &= \sum_{n \geq 0} H_K(\mathcal{F}, n) t^n - 2 \sum_{n \geq 0} H_K(\mathcal{F}, n-1) t^n + \sum_{n \geq 0} H_K(\mathcal{F}, n-2) t^n \\
&= \frac{f(t)}{(1-t)^3} - 2H_K(\mathcal{F}, -1) - 2t \frac{f(t)}{(1-t)^3} + H_K(\mathcal{F}, -2) + tH_K(\mathcal{F}, -1) + t^2 \frac{f(t)}{(1-t)^3} \\
&= \frac{f(t)}{(1-t)} - 2\lambda(R/K) + \lambda(R/K) + t\lambda(R/K) \\
&= \frac{f(t) - (1-t)^2 \lambda(R/K)}{(1-t)}.
\end{aligned}$$

Therefore

$$\sum_{n \geq 0} \Delta^2 [P_K(\mathcal{F}, n) - H_K(\mathcal{F}, n)] t^n = \frac{e_0(\mathcal{F}) - f(t) + (1-t)^2 \lambda(R/K)}{(1-t)} = \sum_{n \geq 0} v_n t^n.$$

Therefore

$$e_0(\mathcal{F}) - f(t) + (1-t)^2 \lambda(R/K) = (1-t) \sum_{n \geq 0} v_n t^n.$$

Hence

$$(1) \quad f(t) = e_0(\mathcal{F}) + (1-t)^2 \lambda(R/K) - (1-t) \sum_{n \geq 0} v_n t^n.$$

Thus,  $f(1) = e_0(\mathcal{F})$ . Differentiating (1) with respect to  $t$ , we get

$$f'(t) = -2(1-t)\lambda(R/K) - (1-t) \sum_{n \geq 0} n v_n t^{n-1} + \sum_{n \geq 0} v_n t^n.$$

Therefore  $g'_1 = f'(1) = \sum_{n \geq 0} v_n$ . Differentiating (1) twice with respect to  $t$ , we get

$$f''(t) = 2\lambda(R/K) - (1-t) \sum_{n \geq 0} n(n-1) v_n t^{n-2} + 2 \sum_{n \geq 0} n v_n t^{n-1}$$

so that

$$g'_2 = f''(1)/2 = \sum_{n \geq 0} n v_n + \lambda(R/K).$$

Therefore

$$g_1 = g'_1 - e_0(\mathcal{F}) = \sum_{n \geq 0} v_n - e_0(\mathcal{F}) = \sum_{n \geq 1} v_n$$

and

$$g_2 = g'_2 - g_1 = \sum_{n \geq 0} n v_n + \lambda(R/K) - \sum_{n \geq 1} v_n = \sum_{n \geq 1} (n-1) v_n + \lambda(R/K).$$

□

**Remark:** The above formulas for  $g_1$  and  $g_2$  generalize the formulas for  $e_1$  and  $e_2$  obtained by Huneke as consequences of his fundamental lemma. To obtain Huneke's formulas for  $e_1$  and  $e_2$ , one simply puts  $K = R$  in the above formulas for  $g_1$  and  $g_2$ .

**Example 3.4.** Let  $k$  be any field and let  $R = k[[x, y]]$ . Let  $I = (x^3, x^2y, y^3)$ . Then  $J = (x^3, y^3)$  is a minimal reduction of  $I$ . Then  $r_J(I) = 3$ ,  $\mathfrak{m}I^n = \mathfrak{m}JI^{n-1}$  for all  $n \geq 2$  and  $\mathfrak{m}I^n : J = \mathfrak{m}I^{n-1}$  for all  $n \geq 1$ . Therefore  $v_n = 0$  for all  $n \geq 2$ . One can also see that  $e_0 = 9$  and  $\lambda(R/\mathfrak{m}I) = 10$ . Hence we have  $v_0 = 9, v_1 = e_0 - \lambda(R/\mathfrak{m}I) + \lambda(R/\mathfrak{m}) = 9 - 10 + 1 = 0$ . Thus  $g'_1 = v_0 + v_1 = 9$  and  $g'_2 = v_1 + \lambda(R/\mathfrak{m}) = 1$  which gives  $g_1 = g'_1 - e_0 = 0$  and  $g_2 = g'_2 - g'_1 + e_0 = 1$ .

The following lemma shows that the behaviour of the superficial elements in  $F_K(\mathcal{F})$  is quite similar to that of superficial elements in  $G(\mathcal{F})$ .

**Lemma 3.5.** *Let  $x$  be a regular element in  $I_1$  such that  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ . Let  $\bar{g}_i$  denote the coefficients of the polynomial corresponding to the function  $\lambda(\bar{R}/\bar{K}\bar{I}_n)$ , where “ $-$ ” denotes modulo( $x$ ). Then  $\bar{g}_i = g_i$  for all  $i = 0, \dots, d-1$ .*

*Proof.* Consider the exact sequence

$$0 \longrightarrow \frac{KI_n : x}{KI_n} \longrightarrow R/KI_n \xrightarrow{x} R/KI_n \longrightarrow R/(KI_n + xR) \longrightarrow 0.$$

Then  $\lambda(\bar{R}/\bar{K}\bar{I}_n) = \lambda(R/(KI_n + xR)) = \lambda(KI_n : x/KI_n)$ . Since  $x^o$  is superficial in  $F_K(\mathcal{F})$  and  $x^*$  is superficial in  $G(\mathcal{F})$ ,  $KI_n : x = KI_{n-1}$  for  $n \gg 0$ , by Lemma 2.3. Hence,  $\lambda(R/(KI_n + xR)) = \lambda(R/KI_n) - \lambda(R/KI_{n-1})$  for  $n \gg 0$ . Therefore

$$\begin{aligned} P_{\bar{K}}(\bar{\mathcal{F}}, n) &= P_K(\mathcal{F}, n) - P_K(\mathcal{F}, n-1) \\ &= \sum_{i=0}^d (-1)^i g_i \binom{n+d-i-1}{d-i} - \sum_{i=0}^d (-1)^i g_i \binom{n+d-i-2}{d-i} \\ &= \sum_{i=0}^{d-1} (-1)^i g_i \binom{n+d-i-2}{d-1-i}. \end{aligned}$$

□

#### 4. COHEN-MACAULAY FIBER CONES

In this section we obtain a lower bound for the Hilbert coefficient  $g_1$ . We will characterize the Cohen-Macaulayness of  $F_K(\mathcal{F})$  in terms of  $g_1$ . In this characterization, We need to assume that  $G(\mathcal{F})$  has almost maximal depth. We will show by an example that we need this assumption for such a characterization. Let  $\gamma(\mathcal{F})$  denote  $\text{grade}_{G(\mathcal{F})_+}(G(\mathcal{F}))$ . We begin by giving a lower bound for the Hilbert coefficient  $g_1$ .

**Proposition 4.1.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d$  with infinite residue field and let  $J \subseteq I_1$  be a minimal reduction of  $I_1$  and let  $K$  be an ideal such that  $I_{n+1} \subseteq KI_n$ . Then  $g_1 \geq \sum_{n \geq 1} \lambda(KI_n + J/J) - \lambda(R/K)$ .*

*Proof.* We apply induction on  $d$ . Let  $d = 1$  and let  $(x) = J$ . For  $i \geq 0$ , from the exact sequence

$$0 \longrightarrow \frac{(KI_{i+1} : x) \cap I_i}{KI_i} \longrightarrow I_i/KI_i \xrightarrow{x} I_{i+1}/KI_{i+1} \longrightarrow I_{i+1}/(KI_{i+1} + xI_i) \longrightarrow 0.$$

it follows that

$$\lambda\left(\frac{I_{i+1}}{KI_{i+1}}\right) - \lambda\left(\frac{I_i}{KI_i}\right) = \lambda\left(\frac{I_{i+1}}{(KI_{i+1} + xI_i)}\right) - \lambda\left(\frac{(KI_{i+1} : x) \cap I_i}{KI_i}\right) \dots\dots (E_i).$$

Summing up  $E_0, E_1, \dots, E_{n-1}$ , we get

$$\lambda(I_n/KI_n) - \lambda(R/K) = \sum_{i=1}^n \lambda(I_i/(KI_i + xI_{i-1})) - \sum_{i=1}^n \lambda((KI_i : x) \cap I_{i-1}/KI_{i-1}).$$

Thus for all  $n \gg 0$ ,

$$f_0 = \lambda(R/K) + \sum_{i \geq 1} \lambda(I_i/(KI_i + xI_{i-1})) - \sum_{i \geq 1} \lambda((KI_i : x) \cap I_{i-1}/KI_{i-1}).$$

By Theorem 4.7 of [HM],  $e_1(\mathcal{F}) = \sum_{i \geq 1} \lambda(I_i/xI_{i-1})$ . Thus

$$\begin{aligned} g_1 &= e_1(\mathcal{F}) - f_0 \\ &= \sum_{i \geq 1} \lambda(I_i/xI_{i-1}) - \lambda(R/K) - \sum_{i \geq 1} \lambda(I_i/(KI_i + xI_{i-1})) + \sum_{i \geq 1} \lambda((KI_i : x) \cap I_{i-1}/KI_{i-1}) \\ &= \sum_{i \geq 1} \lambda((KI_i + xI_{i-1})/xI_{i-1}) + \sum_{i \geq 1} \lambda((KI_i : x) \cap I_{i-1}/KI_{i-1}) - \lambda(R/K). \end{aligned}$$

This implies that  $g_1 \geq \sum_{i \geq 1} \lambda((KI_i + xI_{i-1})/xI_{i-1}) - \lambda(R/K) \geq \sum_{i \geq 1} \lambda(KI_i/J/J) - \lambda(R/K)$ . Hence the result is true for  $d = 1$ . Let us assume that  $d > 1$  and the assertion is true for  $d - 1$ . Choose the generators  $x_1, \dots, x_d$  of  $J$  such that  $x_1^o$  (resp.  $x_1^* \in G(\mathcal{F})$ ) is superficial in  $F_K(\mathcal{F})$  (resp.  $G(\mathcal{F})$ ). Let “ $-$ ” denote images modulo  $(x_1)$ . Then by Lemma 3.5,  $\bar{g}_1 = g_1$ . By induction

$$\begin{aligned} \bar{g}_1 &\geq \sum_{n \geq 1} \lambda(\bar{K}\bar{I}_n + \bar{J}/\bar{J}) - \lambda(\bar{R}/\bar{K}) \\ &= \lambda((KI_n + xR) + (J + xR)/(J + xR)) - \lambda(R/K) \\ &= \sum_{n \geq 1} \lambda(KI_n + J/J) - \lambda(R/K). \end{aligned}$$

□

As a consequence of the above bound, we derive a bound for the multiplicity of the fiber cone given in [CPV, Theorem 2.1].

**Corollary 4.2.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d > 0$  and infinite residue field. Let  $I$  be an  $\mathfrak{m}$ -primary ideal. Then the multiplicity of the fiber cone,  $F(I)$  satisfies*

$$f_0 \leq e_1(I) - e_0(I) + \lambda(R/I) + \mu(I) - d + 1.$$

*Proof.* Since  $f_0 = e_1 - g_1$ , by Proposition 4.1, we get

$$\begin{aligned}
f_0 &\leq e_1 - \sum_{n \geq 1} \lambda(\mathfrak{m}I^n + J/J) + 1 \\
&= e_1 - \lambda(\mathfrak{m}I/\mathfrak{m}J) + 1 - \sum_{n \geq 2} \lambda(\mathfrak{m}I^n + J/J) \\
&= e_1(I) - e_0(I) - d + \lambda(R/I) + \mu(I) + 1 - \sum_{n \geq 2} \lambda(\mathfrak{m}I^n + J/J) \\
&\leq e_1(I) - e_0(I) + \lambda(R/I) + \mu(I) - d + 1.
\end{aligned}$$

□

Now we prove a characterization for Cohen-Macaulayness of the fiber cone in terms of  $g_1$ .

**Theorem 4.3.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d > 0$  with  $R/\mathfrak{m}$  infinite. Let  $\mathcal{F} = \{I_n\}$  be a Hilbert filtration of  $R$  and let  $K$  be an ideal of  $R$  containing  $I_1$ . Let  $J$  be a minimal reduction of  $I_1$ . Assume that  $\gamma(\mathcal{F}) \geq d - 1$ . Then  $F_K(\mathcal{F})$  is Cohen-Macaulay if and only if  $g_1 = \sum_{n \geq 1} \lambda(KI_n + JI_{n-1}/JI_{n-1}) - \lambda(R/K)$ .*

*Proof.* Suppose that  $F_K(\mathcal{F})$  is Cohen-Macaulay. As  $J + KI_1/KI_1$  is generated by a homogeneous system of parameters of degree 1,  $f_0 = \lambda(F_K(\mathcal{F})/JF_K(\mathcal{F})) = \sum_{i \geq 0} \lambda(I_i/(KI_i + JI_{i-1}))$ . Since  $\gamma(\mathcal{F}) \geq d - 1$ , by Theorem 4.7 of [HM],  $e_1 = \sum_{i \geq 1} \lambda(I_i/JI_{i-1})$ . Therefore

$$\begin{aligned}
g_1 &= e_1 - f_0 = \sum_{n \geq 1} \lambda(I_n/JI_{n-1}) - \sum_{n \geq 0} \lambda(I_n/(KI_n + JI_{n-1})) \\
&= \sum_{n \geq 1} \lambda(I_n/JI_{n-1}) - \sum_{n \geq 1} \lambda(I_n/(KI_n + JI_{n-1})) - \lambda(R/K) \\
&= \sum_{n \geq 1} \lambda(KI_n + JI_{n-1}/JI_{n-1}) - \lambda(R/K).
\end{aligned}$$

Conversely, suppose  $g_1 = \sum_{n \geq 1} \lambda(KI_n + JI_{n-1}/JI_{n-1}) - \lambda(R/K)$ . Then by reversing the above steps, one can see that  $f_0 = \sum_{n \geq 0} \lambda(I_n/(KI_n + JI_{n-1})) = \lambda(F_K(\mathcal{F})/JF_K(\mathcal{F}))$ . Therefore  $F_K(\mathcal{F})$  is Cohen-Macaulay. □

The following example shows that the assumption in Theorem 4.3 that  $\text{depth } G(\mathcal{F}) \geq d - 1$  cannot be dropped.

**Example 4.4.** Let  $R = k[[x, y]]$ ,  $\mathfrak{m} = (x, y)$  and  $I = (x^4, x^3y, xy^3, y^4)$ . Then  $J = (x^4, y^4)$  is a minimal reduction of  $I$  and  $I^3 = JI^2$ . Note that  $I^n = \mathfrak{m}^{4n}$  for all  $n \geq 2$ . We compute the Hilbert coefficients of  $I$ . Since  $I^n = \mathfrak{m}^{4n}$  for all  $n \geq 2$ ,  $\lambda(R/I^n) = \lambda(R/\mathfrak{m}^{4n}) = \binom{4n+1}{2} = e_0(I) \binom{n+1}{2} - e_1(I)n + e_2(I)$ . Solving the equation by putting various values for  $n$ , we get  $e_0(I) = 16$ ,  $e_1(I) = 6$ ,  $e_2(I) = 0$ . From direct computations one can see  $\lambda(I/J) = 5$  and  $\lambda(I^2/JI) = 2$ . Hence  $e_1(I) < \sum_{n \geq 1} \lambda(I^n/JI^{n-1})$ . Therefore  $\text{depth } G(I) = 0$ . Since  $I^n = \mathfrak{m}^{4n}$  for all  $n \geq 2$ ,  $\mu(I^n) = \lambda(\mathfrak{m}^{4n}/\mathfrak{m}^{4n+1}) = \binom{4n+1}{1}$  for all  $n \geq 1$ . Therefore  $f_0 = 4$  so that  $g_1 = e_1 - f_0 = 2$ . Also, one can see that  $\lambda(\mathfrak{m}I + J/J) \neq 0$  and

$\mathfrak{m}I^n \subseteq JI^{n-1}$  for all  $n \geq 2$ . Then  $\lambda(\mathfrak{m}I + J/J) = \lambda(\mathfrak{m}I/\mathfrak{m}I \cap J) = \lambda(\mathfrak{m}I/\mathfrak{m}J) = 3$ . Therefore  $\sum_{n \geq 1} \lambda(\mathfrak{m}I^n + JI^{n-1}/JI^{n-1}) - 1 = 2 = g_1$ . The Hilbert Series of the fiber cone is given by

$$H(F(I), t) = \frac{1 + 2t + 2t^2 - t^3}{(1 - t)^2}.$$

Since the numerator contains a negative coefficient,  $F(I)$  is not Cohen-Macaulay.

## 5. FIBER CONES WITH ALMOST MAXIMAL DEPTH

In this section we present a characterization for the fiber cone  $F_K(\mathcal{F})$  to have almost maximal depth in terms of  $g_1$ . This is an analogue of the Huckaba-Marley characterization for the associated graded ring to have almost maximal depth referred above. We will need to invoke a result due to Cortadellas and Zarzuela from [CZ] which gives a criterion for a sequence of degree one elements in  $F_K(\mathcal{F})$  to be a regular sequence. We take this opportunity to present a simple proof of their result since it is a very basic result for detecting regular sequences in fiber cones.

**Theorem 5.1.** *Let  $(R, \mathfrak{m})$  be a Noetherian local ring,  $\mathcal{F}$  a filtration of ideals,  $K$  an ideal containing  $I_1$  and  $x_1, \dots, x_k \in I_1$ . Assume that*

- (i)  $x_1, \dots, x_k$  is a regular sequence in  $R$ .
- (ii)  $x_1^* \dots, x_k^* \in G(\mathcal{F})$  is a regular sequence.
- (iii)  $x_1^o, \dots, x_k^o \in F_K(\mathcal{F})$  is a superficial sequence.

*Then  $\text{depth}_{(x_1^o, \dots, x_k^o)} F_K(\mathcal{F}) = k$  if and only if  $(x_1, \dots, x_k) \cap KI_n = (x_1, \dots, x_k)KI_{n-1}$  for all  $n \geq 1$ .*

*Proof.* We induct on  $k$ . Let  $k = 1$ . Let  $(x) \cap KI_n = xKI_{n-1}$  for all  $n \geq 1$ . Then  $KI_n : x = KI_{n-1}$  for all  $n \geq 1$  and hence  $x^o$  is regular in  $F_K(\mathcal{F})$ . Suppose  $x^o$  is regular in  $F_K(\mathcal{F})$ . Let  $n \geq 1$  and  $y \in (x) \cap KI_n$ . Let  $y = rx$  for some  $r \in R$ . Then  $r \in KI_n : x \subseteq I_n : x$ . Since  $x^*$  is a nonzerodivisor in  $G(\mathcal{F})$ ,  $I_n : x = I_{n-1}$  for all  $n \geq 1$ . Therefore  $r \in I_{n-1}$ . Hence  $r \in (KI_n : x) \cap I_{n-1} = KI_{n-1}$  so that  $y \in xKI_{n-1}$ . Now assume that  $k > 1$  and the result is true for all  $l \leq k - 1$ . Put  $J = (x_1, \dots, x_k)$ ,  $J^o = (x_1^o, \dots, x_k^o) \subseteq F_K(\mathcal{F})$  and  $J^* = (x_1^*, \dots, x_k^*) \subseteq G(\mathcal{F})$ . Let “ $-$ ” denote images modulo  $(x_1)$ . Then  $F_K(\mathcal{F})/x_1^o F_K(\mathcal{F}) \cong F_{\bar{K}}(\bar{\mathcal{F}})$ . Assume  $J \cap KI_n = JKI_{n-1}$  for all  $n \geq 1$ . Then

$$\bar{J} \cap \bar{K} \bar{I}_n = (J + x_1 R) \cap (KI_n + x_1 R) = J \cap (KI_n + x_1 R) = (J \cap KI_n) + x_1 R = JKI_{n-1} + x_1 R = \bar{J} \bar{K} \bar{I}_{n-1}.$$

By induction  $\text{depth}_{\bar{J}^o} F_{\bar{K}}(\bar{\mathcal{F}}) = k - 1$ . Thus  $x_1^o$  is regular in  $F_K(\mathcal{F})$  and hence,  $\text{depth}_{J^o}(F_K(\mathcal{F})) = k$ , by Lemma 2.7. Conversely assume that  $\text{depth}_{J^o}(F_K(\mathcal{F})) = k$ . Since  $x_1^o$  is superficial in  $F_K(\mathcal{F})$  and  $\text{depth} F_K(\mathcal{F}) > 0$ ,  $x_1^o$  is regular in  $F_K(\mathcal{F})$ . Then  $\text{depth}_{\bar{J}^o}(F(\bar{I})) = k - 1$ . Hence, by induction,  $\bar{J} \cap \bar{K} \bar{I}_n = \bar{J} \bar{K} \bar{I}_{n-1}$ . Therefore,  $J \cap KI_n + x_1 R = JKI_{n-1} + x_1 R$ . Hence

$$\begin{aligned} J \cap KI_n &= JKI_{n-1} + (x_1 R \cap (J \cap KI_n)) \\ &= JKI_{n-1} + (x_1 R \cap KI_n) \\ &= JKI_{n-1} + (x_1 KI_{n-1}) = JKI_{n-1}. \end{aligned}$$

Therefore  $J \cap KI_n = JKI_{n-1}$ . □

We need the following lemma in the proof of the characterization for the fiber cone to have depth at least  $d - 1$ .

**Lemma 5.2.** *Let  $(R, \mathfrak{m})$  be a  $d$ -dimensional Cohen-Macaulay local ring and let  $\mathcal{F} = \{I_n\}$  be a Hilbert filtration of  $R$  such that  $\gamma(\mathcal{F}) \geq d - 1$ . Let  $K$  be an ideal of  $R$  containing  $I_1$ . Let  $J = (x_1, \dots, x_d)$  be a minimal reduction of  $I_1$  such that  $x_1^*, \dots, x_{d-1}^*$  is a regular sequence in  $G(\mathcal{F})$ . If  $KI_n \cap (x_1, \dots, x_{d-1}) \subseteq JKI_{n-1}$  for all  $n \geq 1$ , then  $x_1^o, \dots, x_{d-1}^o$  is a regular sequence in  $F_K(\mathcal{F})$ .*

*Proof.* Since  $x_1^*, \dots, x_{d-1}^*$  is a regular sequence in  $G(\mathcal{F})$ , by Theorem 5.1, it is enough to show that  $KI_n \cap (x_1, \dots, x_{d-1}) = (x_1, \dots, x_{d-1})KI_{n-1}$  for all  $n \geq 1$ . Induct on  $n$ . Let  $z \in KI_1 \cap (x_1, \dots, x_{d-1})$ . Write  $z = \sum_{i=1}^{d-1} r_i x_i$  for  $r_i \in R$ . Since  $KI_1 \cap (x_1, \dots, x_{d-1}) \subseteq (x_1, \dots, x_d)K$ ,  $z = \sum_{i=1}^{d-1} s_i x_i + px_d$  for some  $p, s_i \in K$ . Then  $px_d \in (x_1, \dots, x_{d-1})$  and hence  $p \in (x_1, \dots, x_{d-1})$ . Since  $x_d \in K$ ,  $z \in (x_1, \dots, x_{d-1})K$ . Now assume that  $n \geq 2$  and for all  $l < n$ ,

$$KI_l \cap (x_1, \dots, x_{d-1}) = (x_1, \dots, x_{d-1})KI_{l-1}.$$

Let  $z \in KI_n \cap (x_1, \dots, x_{d-1}) \subseteq JKI_{n-1}$ . Write  $z = \sum_{i=1}^{d-1} r_i x_i = \sum_{i=1}^{d-1} s_i x_i + px_d$ , where  $r_i \in R$ ,  $p, s_i \in KI_{n-1}$ . Then  $px_d \in (x_1, \dots, x_{d-1})$  and hence  $p \in (x_1, \dots, x_{d-1})$ . Therefore

$$KI_n \cap (x_1, \dots, x_{d-1}) = (x_1, \dots, x_{d-1})KI_{n-1} + x_d(KI_{n-1} \cap (x_1, \dots, x_{d-1})).$$

By induction  $KI_{n-1} \cap (x_1, \dots, x_{d-1}) \subseteq (x_1, \dots, x_{d-1})KI_{n-2}$ . Hence  $p \in (x_1, \dots, x_{d-1})KI_{n-2}$  so that  $px_d \in (x_1, \dots, x_{d-1})KI_{n-1}$ . Therefore  $z \in (x_1, \dots, x_{d-1})KI_{n-1}$ .  $\square$

We prove a necessary and sufficient condition for the fiber cone to have depth at least  $d - 1$  in terms of  $g_1$  in the following theorem.

**Theorem 5.3.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d > 0$ . Let  $\mathcal{F}$  be a Hilbert filtration,  $K$  an ideal such that  $I_{n+1} \subseteq KI_n$  for all  $n \geq 0$  and  $J$  a minimal reduction of  $I_1$ . Assume that  $\gamma(\mathcal{F}) \geq d - 1$ . Then  $g_1 = \sum_{n \geq 1} \lambda(KI_n/JKI_{n-1}) - \lambda(R/K)$  if and only if  $\text{depth } F_K(\mathcal{F}) \geq d - 1$ .*

*Proof.* We induct on  $d$ . Let  $d = 1$ . In this case, we need to show that  $g_1$  is equal to the summation given in the statement of the theorem. Let  $(x)$  be a reduction of  $I_1$ . From the proof of Proposition 4.1 we get,

$$g_1 = \sum_{n \geq 1} \lambda((KI_n + xI_{n-1})/xI_{n-1}) + \sum_{n \geq 1} \lambda((KI_n : x) \cap I_{n-1}/KI_{n-1}) - \lambda(R/K).$$

*Claim :* For all  $n \geq 1$ ,  $(KI_n : x) \cap I_{n-1}/KI_{n-1} \cong xI_{n-1} \cap KI_n/xKI_{n-1}$ .

Consider the multiplication map  $\mu_x : (KI_n : x) \cap I_{n-1}/KI_{n-1} \longrightarrow xI_{n-1} \cap KI_n/xKI_{n-1}$ . Let  $y = xs \in xI_{n-1} \cap KI_n$  for some  $s \in I_{n-1}$ . Then  $s \in (KI_n : x) \cap I_{n-1}$ . Therefore,  $\mu_x$  is surjective. Let  $y \in (KI_n : x) \cap I_{n-1}$  and  $xy \in xKI_{n-1}$ . Since  $x$  is regular in  $R$ ,  $y \in KI_{n-1}$ . Hence  $\mu_x$  is injective so that  $\mu_x$  is an isomorphism. Therefore we have,

$$\begin{aligned} g_1 &= \sum_{n \geq 1} [\lambda(KI_n/(KI_n \cap xI_{n-1})) + \lambda((xI_{n-1} \cap KI_n)/xKI_{n-1})] - \lambda(R/K) \\ &= \sum_{n \geq 1} \lambda(KI_n/xKI_{n-1}) - \lambda(R/K). \end{aligned}$$

Assume now that  $d > 1$ . Choose  $x_1, \dots, x_d$  such that  $J = (x_1, \dots, x_d)$ ,  $x_1^*, \dots, x_d^*$  is a superficial sequence in  $G(\mathcal{F})$  and  $(x_1^o, \dots, x_d^o)$  is a superficial sequence in  $F_K(\mathcal{F})$ . Since  $\gamma(\mathcal{F}) \geq d-1$ ,  $x_1^*, \dots, x_{d-1}^*$  is a regular sequence in  $G(\mathcal{F})$  (existence of such a generating set can be derived from Proposition A.2.4 of [Ma]). Suppose  $g_1 = \sum_{n \geq 1} \lambda(KI_n/JKI_{n-1}) - \lambda(R/K)$ . Let “ $-$ ” denote images modulo  $(x_1, \dots, x_{d-1})$ . Then by Lemma 3.5,  $g_1 = \bar{g}_1$  and

$$\begin{aligned} \bar{g}_1 &= \sum_{n \geq 1} \lambda(\bar{K}\bar{I}_n/\bar{J}\bar{K}\bar{I}_{n-1}) - \lambda(\bar{R}/\bar{K}) \quad (\text{Since } \dim \bar{R} = 1) \\ &= \sum_{n \geq 1} \lambda((KI_n + (x_1, \dots, x_{d-1})) / (JKI_{n-1} + (x_1, \dots, x_{d-1}))) - \lambda(R/K) \\ &= \sum_{n \geq 1} \lambda(KI_n / (JKI_{n-1} + KI_n \cap (x_1, \dots, x_{d-1}))) - \lambda(R/K). \end{aligned}$$

By assumption  $g_1 = \sum_{n \geq 1} \lambda(KI_n/JKI_{n-1}) - \lambda(R/K)$ . Therefore,  $(x_1, \dots, x_{d-1}) \cap KI_n \subseteq JKI_{n-1}$ . Hence by Lemma 5.2,  $x_1^o, \dots, x_{d-1}^o$  is a regular sequence in  $F_K(\mathcal{F})$ .

Conversely, let  $\text{depth } F_K(\mathcal{F}) \geq d-1$ . Choose  $x_1 \in I_1$  such that  $x_1^*$  is regular in  $G(\mathcal{F})$  and  $x_1^o$  is regular in  $F_K(\mathcal{F})$ . Hence  $F_K(\mathcal{F})/x_1^o F_K(\mathcal{F}) \cong F_{\bar{K}}(\bar{\mathcal{F}})$  and  $g_1 = \bar{g}_1$ , where “ $-$ ” denote images modulo  $(x_1)$ . Then  $\text{depth } F_K(\mathcal{F})/x_1^o F_K(\mathcal{F}) \geq d-2$ . By induction

$$\begin{aligned} \bar{g}_1 &= \sum_{n \geq 1} \lambda(KI_n + x_1 R / JKI_{n-1} + x_1 R) - \lambda(R/K) \\ &= \sum_{n \geq 1} \lambda(KI_n / (JKI_{n-1} + (x_1 R \cap KI_n))) - \lambda(R/K). \end{aligned}$$

Since  $x_1^o$  is regular in  $F_K(\mathcal{F})$  and  $x_1^*$  is regular in  $G(\mathcal{F})$ ,  $x_1 R \cap KI_n = x_1 KI_{n-1}$ . Therefore

$$\bar{g}_1 = \sum_{n \geq 1} \lambda(KI_n / (JKI_{n-1} + x_1 KI_{n-1})) - \lambda(R/K) = \sum_{n \geq 1} \lambda(KI_n / JKI_{n-1}) - \lambda(R/K).$$

Since  $g_1 = \bar{g}_1$ , the assertion follows.  $\square$

**Remark 5.4.** From the above proof we obtain an upper bound for the coefficient  $g_1$ :

$$g_1 \leq \lambda(KI_n / KJI_{n-1}) - \lambda(R/K).$$

## 6. COHEN-MACAULAY FIBER CONES OF IDEALS WITH MINIMAL MULTIPLICITY

Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d > 0$ . Let  $I$  be an  $\mathfrak{m}$ -primary ideal of  $R$  and  $J$  be a minimal reduction of  $I$ . Let  $K$  be an ideal containing  $I$ . Let  $F_K(I)$  be the fiber cone of  $I$  with respect to  $K$  and let  $G(I)$  be the associated graded ring of  $I$ . Let  $\mu(I)$  denote the minimum number of generators of  $I$ . It is known that  $e_0(I) + d - \lambda(R/I) \geq \mu(I)$ , [G]. Goto defined an ideal  $I$  to have minimal multiplicity if  $e_0(I) + d - \lambda(R/I) = \mu(I)$ . It can be seen that  $I$  has minimal multiplicity if and only if for any minimal reduction  $J$  of  $I$ ,  $I\mathfrak{m} = J\mathfrak{m}$ . We generalize this notion. An ideal  $I$  is said to have minimal multiplicity with respect to  $K$  if  $KI = KJ$  for any minimal reduction  $J$  of  $I$ . Let  $H_K(I, n) = \lambda(R/KI^n)$  and  $P_K(I, n) = \sum_{i=0}^d (-1)^i g_i \binom{n+d-i-1}{d-i}$  be the corresponding polynomial.

**Proposition 6.1.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d$ .*

- (i) *If  $I$  has minimal multiplicity with respect to  $K$ , then  $g_1 = -\lambda(R/K)$ .*
- (ii) *If  $KI \cap J = KJ$  and  $g_1 = -\lambda(R/K)$  then  $I$  has minimal multiplicity.*
- (iii) *The ideal  $I$  has minimal multiplicity if and only if  $g_1 = -1$ .*

*Proof.* (i) Suppose  $I$  has minimal multiplicity with respect to  $K$ . Let  $J$  be a minimal reduction of  $I$ . Then  $KI^n = KJ^n$  for all  $n \geq 1$ . Therefore for all  $n \geq 1$ ,

$$\begin{aligned} \lambda(R/KI^n) &= \lambda(R/KJ^n) = \lambda(R/J^n) + \lambda(J^n/KJ^n) \\ &= e_0(I) \binom{n+d-1}{d} + \lambda(R/K) \binom{n+d-1}{d-1}. \end{aligned}$$

Hence  $g_1 = -\lambda(R/K)$ .

- (ii) Suppose  $g_1 = -\lambda(R/K)$  and  $KI \cap J = KJ$ . By Proposition 4.1,  $g_1 \geq \sum_{n \geq 1} \lambda(KI^n + J/J) - \lambda(R/K)$ . Hence  $KI \subseteq J$ . Thus  $KI = KI \cap J = KJ$ . Hence  $I$  has minimal multiplicity with respect to  $K$ .
- (iii) Follows from (i) and (ii) since  $\mathfrak{m}I \cap J = \mathfrak{m}J$ . □

In the following Proposition we generalize, the equivalence of Cohen-Macaulayness of the fiber cone and the almost maximal depth for  $G(I)$ , given in Proposition 2.5 of [G].

**Proposition 6.2.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring,  $I$  an  $\mathfrak{m}$ -primary ideal,  $J$  a minimal reduction of  $I$  and  $K$  an ideal containing  $I$ . Suppose that there exists an integer  $t$  such that*

$$KI^n \cap J = KJI^{n-1} \text{ for } n = 1, \dots, t \quad \text{and} \quad KI^{t+1} = KJI^t. \quad (*)$$

*Then  $F_K(I)$  is Cohen-Macaulay if and only if  $\gamma(I) \geq d - 1$ .*

*Proof.* From  $(*)$ , it follows that  $KI^n \cap J = KI^n \cap JI^{n-1} = KJI^{n-1}$  for all  $n = 1, \dots, t$ . Therefore, from Remark 5.4 and Proposition 4.1, it follows that

$$g_1 = \sum_{n \geq 1} \lambda(KI^n/KJI^{n-1}) - \lambda(R/K) = \sum_{n \geq 1} \lambda(KI^n + JI^{n-1}/JI^{n-1}) - \lambda(R/K).$$

Therefore by Theorem 4.3, if  $\gamma(I) \geq d - 1$ , then  $F_K(I)$  is Cohen-Macaulay.

Assume that  $F_K(I)$  is Cohen-Macaulay. Then  $J + KI/KI$  is system of parameters for  $F_K(I)$  and

$$f_0 = \lambda(F_K(I)/JF_K(I)) = \sum_{n \geq 0} \lambda(I^n/KI^n + JI^{n-1}) = \lambda(R/K) + \sum_{n \geq 1} \lambda(I^n/KI^n + JI^{n-1}).$$

Hence

$$\begin{aligned} e_1(I) &= f_0 + g_1 \\ &= \lambda(R/K) + \sum_{n \geq 1} \lambda(I^n/KI^n + JI^{n-1}) + \sum_{n \geq 1} \lambda(KI^n + JI^{n-1}/JI^{n-1}) - \lambda(R/K) \\ &= \sum_{n \geq 1} \lambda(I^n/JI^{n-1}). \end{aligned}$$

Therefore, by Theorem 4.7 of [HM],  $\gamma(I) \geq d - 1$ . □

**Corollary 6.3.** *Let  $(R, \mathfrak{m})$  be a  $d$ -dimensional Cohen-Macaulay local ring and let  $I$  be an  $\mathfrak{m}$ -primary ideal with minimal multiplicity with respect to  $K \supseteq I$ . Suppose  $KI \cap J = KJ$ . Then  $F_K(I)$  is Cohen-Macaulay if and only if  $\gamma(I) \geq d - 1$ .*

**Theorem 6.4.** *Let  $(R, \mathfrak{m})$  be a Cohen-Macaulay local ring of dimension  $d$  and let  $I$  be an  $\mathfrak{m}$ -primary ideal of  $R$  with minimal multiplicity. Then the following statements are equivalent:*

1.  $G(I)$  is Cohen-Macaulay.
2.  $F(I)$  is Cohen-Macaulay and  $r(I) \leq 1$ .
3.  $r(I) \leq 1$ .

*Proof.* (1)  $\Rightarrow$  (2) : Since  $I$  has minimal multiplicity,  $I\mathfrak{m} = J\mathfrak{m}$  for any minimal reduction  $J$  of  $I$ . Assume that  $G(I)$  is Cohen-Macaulay. Then by Proposition 6.3,  $F(I)$  is Cohen-Macaulay. Therefore

$$\begin{aligned}
 f_0 &= \sum_{n \geq 0} \lambda(I^n/JI^{n-1} + \mathfrak{m}I^n) = 1 + \sum_{n \geq 1} \lambda(I^n/JI^{n-1}) \quad (\text{since } \mathfrak{m}I^n \subseteq JI^{n-1} \text{ for all } n \geq 1) \\
 &= 1 + \lambda(I/J) + \sum_{n \geq 2} \lambda(I^n/JI^{n-1}) \\
 &= 1 + e_0(I) - \lambda(R/I) + \sum_{n \geq 2} \lambda(I^n/JI^{n-1}) \\
 &= 1 + \mu(I) - d + \sum_{n \geq 2} \lambda(I^n/JI^{n-1})
 \end{aligned}$$

Since  $I^2 \subseteq I\mathfrak{m} = J\mathfrak{m} \subseteq J$ ,  $I^2 = I^2 \cap J = IJ$ , as  $G(I)$  is Cohen-Macaulay.

(2)  $\Rightarrow$  (3) : Clear.

(3)  $\Rightarrow$  (1) : This is known but we recall the argument. Assume that  $r(I) \leq 1$ . Then  $I^{n+1} = JI^n$  for all  $n \geq 1$  so that  $JI^n = I^{n+1} \cap J$  for all  $n \geq 1$ . Therefore by [VV],  $x_1^*, \dots, x_d^*$  is a regular sequence in  $G(I)$ . Hence  $G(I)$  is Cohen-Macaulay.  $\square$

**Example 6.5.** Let  $R = k[[t^4, t^5, t^6, t^7]]$  and let  $I = (t^4, t^5, t^6)$ . Then  $J = (t^4)$  is a minimal reduction of  $I$ . It can easily be checked that  $I\mathfrak{m} = J\mathfrak{m}$ . Hence  $F(I)$  is Cohen-Macaulay. Since  $I$  has minimal multiplicity,  $g_1 = -1$ . Since  $\mathfrak{m}I^n \subseteq J$  for all  $n \geq 1$ ,  $g_1 = -1 = \sum_{n \geq 1} \lambda(\mathfrak{m}I^n + JI^{n-1}/JI^{n-1}) - \lambda(R/\mathfrak{m})$ .

## REFERENCES

- [BH] W. Bruns and J. Herzog, *Cohen-Macaulay rings*, Revised Edition, Cambridge Studies in Advanced Mathematics, 39. Cambridge University Press, Cambridge, 1998
- [C] A. Corso, *Sally modules of  $\mathfrak{m}$ -primary ideals in local rings*, preprint, arXiv:math.AC/0309027.
- [CGPU] A. Corso, L. Ghezzi, C. Polini and B. Ulrich, *Cohen-Macaulayness of special fiber rings* Comm. Algebra 31 (special issue in honor of S. Kleiman) (2003), 3713-3734.
- [CPV] A. Corso, C. Polini and W. V. Vasconcelos, *Multiplicity of the special fiber ring of blowups*, preprint, arXiv:math.AC/0307037
- [CZ] T. Cortadellas and S. Zarzuela, *On the depth of the fiber cone of filtrations*, J. Algebra **198** (1997), no. 2, 428-445.
- [DRV] C. D'Cruz, K. N. Raghavan and J. K. Verma, *Cohen-Macaulay fiber cones*, Commutative Algebra, Algebraic Geometry and Computational Methods (Hanoi, 1996), 233-246, Springer, Singapore, 1999

- [DV] C. D'Cruz and J. K. Verma, *Hilbert series of fiber cones of almost minimal mixed multiplicity*, J. Algebra **251** (2002), no. 1, 98–109.
- [G] S. Goto, *Cohen-Macaulayness and negativity of  $A$ -invariants in Rees algebras associated to  $\mathfrak{m}$ -primary ideals of minimal multiplicity*, Commutative algebra, homological algebra and representation theory (Catania/Genoa/Rome, 1998). J. Pure Appl. Algebra **152** (2000), no. 1-3, 93–107.
- [HM] S. Huckaba and T. Marley, *Hilbert coefficients and depth of associated graded rings*, J. London Math. Soc. (2) **56** (1997), 64–76.
- [H] C. Huneke, *Hilbert functions and symbolic powers*, Michigan Math. J. **34** (1987), no. 2, 293–318.
- [HH] R. Hübl and C. Huneke, *Fiber cones and the integral closure of ideals*, Collect. Math. **52** (2001), no. 1, 85–100.
- [Ma] T. Marley, *Hilbert functions of ideals in Cohen-Macaulay rings*, Thesis, Purdue University (1989).
- [S1] J. D. Sally, *On the associated graded ring of a local Cohen-Macaulay ring*, J. Math. Kyoto Univ., **17**(1977), 19–21.
- [S2] J. D. Sally, *Cohen-Macaulay local rings of embedding dimension  $e + d - 2$* , J. Algebra **83** (1983) 393–408.
- [VV] P. Valabrega and G. Valla, *Form rings and regular sequences*, Nagoya Math. J. **72** (1978), 93–101.

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, COLABA, MUMBAI, INDIA - 400005

*E-mail address:* `jayan@math.tifr.res.in`

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY BOMBAY, POWAI, MUMBAI, INDIA - 400076

*E-mail address:* `jkv@math.iitb.ac.in`